

Cooling of a Λ -type three-level atom in a high finesse optical cavity

Lei Tan^{1,2, a}, Li-Wei Liu¹, Yan-Fen Sun¹

¹Institute of Theoretical Physics, Lanzhou University, Lanzhou 730000, China

²Key Laboratory for Magnetism and Magnetic materials of the Ministry of Education, Lanzhou University, Lanzhou 730000, China

Received: date / Revised version: date

Abstract. A theoretical study is carried out for the cavity cooling of a Λ -type three level atom in a high-finesse optical cavity with a weakly driven field. Analytical expressions for the friction, diffusion coefficients and the equilibrium temperatures are obtained by using the Heisenberg equations, then they are calculated numerically and shown graphically as a function of controlling parameters. For a suitable choice of these parameters, the dynamics of the cavity field interaction with the Λ -type three-level atom introduces a sisypus cooling mechanism yielding lower temperatures below the Doppler limit and allowing larger cooling rate, avoiding the problems induced by spontaneous emission.

PACS. 3 2.80.Pj; 42.50.Vk.

1 Introduction

The technical developments of laser cooling and trapping greatly promote atom manipulation, and have opened a new period in cavity QED experimentation. New cavity cooling mechanism[1,2], has been proposed for a single atom strongly coupled to a high-finesse optical cavity, and the cooling mechanism results in an extended storage time and improved localization of atoms. An estimate shows that the observed cooling rate is at least five times large than the cooling rate which can be achieved by free-space cooling methods for comparable excitation of the atom[2]. Compared to all conventional laser cooling methods[3,4,5], cavity cooling does not rely on repeated cycles of optical pumping and spontaneous emission of photons by the atom, but on the escape of photons from the cavity. Moreover, as there is no requirement in cavity cooling for a closed multilevel system, it is an attractive approach for creating ultracold molecules[6, 7]. Recently, cold atoms and molecules[8,9,10,11] are now the basis of many new areas of fundamental physics and technology, and are the central to investigation of the Bose-Einstein condensates[12,13], and quantum information processing[14], etc.

A single atom coupled to a single mode of an optical cavity is the archetype model of dissipative electrodynamics. As is well known, the atom-cavity system has two important characteristics, the first is the cavity mirrors confining the light, which can lead to significantly modified of atom by the cavity field. On the contrary, the

back action of the atom on the intra-cavity field is also notable; The second is the cavity mirrors offering an extra loss channel, which is responsible for a new dissipative and can efficiently damp atomic motion[1,2,15,16]. It is well known that the cavity cooling depends on the optical forces, as well as on the dissipation properties. This motivates us to investigate these forces and dissipation properties in detail. These forces are quite substantial, which can be viewed as the dipole force and friction force. However, dissipation inevitably leads to momentum diffusion, which will heat the atom. In our paper, there are two major contributions to the momentum diffusion. One is the random momentum transfer of absorbed and emitted photons, the second referred to the momentum diffusion is due to the fluctuations of the dipole force. Finally, we can obtain the equilibrium temperatures when the contributions of the friction and heating cancel.

Even though a two-level atom model is usually used to analyze the cavity cooling in a high-finesse optical cavity[17, 18,19,20,21], the nature of a three-level atom-cavity system is quantitatively different from that of a two-level atom system. For a there-level atom in the optical cavity, there are more controlling parameters, namely, two detunings, two Rabi frequencies, cavity detunings, etc. Moreover, the extended level structure also provides flexibility. Because of these reasons, it is of great interest to extend the investigation on the optical forces and momentum diffusion on the three-level atom in a high-finesse optical cavity. Very recently, we have studied the friction force of a V-type three-level atoms in a high-Q cavity [22]. Here, we will concentrate our attention on the variation of the

^a Corresponding author, *Present address:* tanlei@lzu.edu.cn




Fig. 1. Configuration of a Λ -type three-level atom coupled to a single-mode cavity and driven by a laser field.

friction, diffusion coefficients and the equilibrium temperature of a Λ -type three-level atom in high-finesse cavity, then the full analytical expressions of the friction, diffusion coefficients and the equilibrium temperature are obtained. For a suitable choice of the parameters, we can obtain lower equilibrium temperatures than the Doppler limit. For simplicity, we will only consider one-dimensional situation.

The paper is organized as follows. In Section 2 we will give the full analytical solutions for the optical forces and friction coefficient by using Heisenberg equations. According to quantum regression theorem we obtain the momentum diffusion and get the equilibrium temperature in Section 3. Section 4 we study the variations of the friction, diffusion coefficients and the equilibrium temperature with detunings, coupling strengths, etc. The friction, diffusion coefficients and the equilibrium temperature are calculated numerically and shown graphically as a function of controlling parameters. Some concluding remarks are presented in Section 5.

2 Dipole force and friction force in the Λ -type three-level atom

Consider a Λ -type three-level atom strongly coupled to a single mode of the electromagnetic field contained in a high-finesse optical cavity with the cavity photons decay at a rate κ . The schematic sketch of the basic system is given in Fig. 1. The upper level is $|3\rangle$ (energy 0), while the two lower levels are $|2\rangle$ (energy $-\hbar\omega_{32}$), and $|1\rangle$ (energy $-\hbar\omega_{31}$). The transition between $|3\rangle$ and level $|2\rangle$ ($|1\rangle$) is mediated by frequency ω_{32} (ω_{31}). The coupling strengths are denoted as $g_1(x) = g_0 \cos(k_1 x)$ and $g_2(x) = g_0 \cos(k_2 x)$, respectively, k_1, k_2 are the wave vector. The transition between states $|2\rangle$ and $|1\rangle$ is forbidden. The state $|3\rangle$ decays radiatively into $|1\rangle$ ($|2\rangle$) at rate Γ_1 (Γ_2). In general, the system is pumped along the cavity axis by a coherent laser field of frequency ω_p and effective amplitude η . The master equation describing the model in Fig. 1 is given in the rotating-wave approximation and in a frame rotating with the pump frequency ω_p as[20]

$$\dot{\rho} = -\frac{i}{\hbar}[H_{ac}, \rho] - \frac{i}{\hbar}[H_p, \rho] + L_\kappa \rho + L_\gamma \rho, \quad (1)$$

where

$$\begin{aligned} H_{ac} &= -\hbar\Delta_c a^\dagger a + \hbar\Delta_1 A_{11} + \hbar\Delta_2 A_{22} \\ &\quad + \hbar g_1(x)(a^\dagger A_{13} + A_{31}a) + \hbar g_2(x)(a^\dagger A_{23} + A_{32}a), \\ H_p &= i\hbar\eta(a^\dagger - a), \\ L_\kappa \rho &= \kappa(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a), \\ L_\gamma \rho &= \Gamma_1(2A_{13}\rho A_{31} - \rho A_{31}A_{13} - A_{31}A_{13}\rho) \\ &\quad + \Gamma_2(2A_{23}\rho A_{32} - \rho A_{32}A_{23} - A_{32}A_{23}\rho) \end{aligned} \quad (2)$$

with the atomic detunings $\Delta_1 = \omega_p - \omega_{31}$ and $\Delta_2 = \omega_p - \omega_{32}$, and cavity detuning $\Delta_c = \omega_p - \omega_c$. The term H_{ac} represents the energy of the atomic system and the interaction of the atom with the cavity field, the term H_p is the pump hamiltonian. $L_\kappa \rho$ and $L_\gamma \rho$ describe the decay of the resonator mode and the atomic damping to background modes, respectively. Instead of solving the master equation one can equivalently deal with the Heisenberg equations for atomic and cavity field mode operators. Considering the spontaneous emission and cavity decay, the Heisenberg equations about the atomic coupling to the vacuum field can be obtained[23].

$$\dot{a} = i\Delta_c a - ig_1(x)A_{13} - ig_2(x)A_{23} - \kappa a + \eta, \quad (3)$$

$$\dot{A}_{13} = i\Delta_1 A_{13} + ig_1(x)(A_{33} - A_{11})a - \Gamma_1 A_{13}, \quad (4)$$

$$\dot{A}_{23} = i\Delta_2 A_{23} + ig_2(x)(A_{33} - A_{22})a - \Gamma_2 A_{23}, \quad (5)$$

where the atomic operator $A_{ij} = |i\rangle\langle j|$ satisfies the commutator relation $[A_{ij}, A_{kl}] = A_{il}\delta_{jk} - A_{kj}\delta_{il}$. In order to linearize Eq. (3) to Eq. (5), we assume that the cavity is very weakly driven then the approximation of small saturation and weak excitation of an atom can be fitted[24, 25]. In this case there is at most one photon in the cavity and we assume that $A_{11} + A_{22} + A_{33} = 1$. For simplicity, we neglect the population of the excited state and assume $\langle A_{11} \rangle = F$ and $\langle A_{22} \rangle = 1 - F$, Thus,

$$\begin{aligned} \langle (A_{33} - A_{11})a \rangle &= -F\langle a \rangle, \\ \langle (A_{33} - A_{22})a \rangle &= -(1 - F)\langle a \rangle. \end{aligned} \quad (6)$$

Then the time evolution of the expectation values can be written as

$$\begin{aligned} \langle \dot{a} \rangle &= i\Delta_c \langle a \rangle - ig_1(x)\langle A_{13} \rangle - ig_2(x)\langle A_{23} \rangle - \kappa \langle a \rangle + \eta, \\ \langle \dot{A}_{13} \rangle &= i\Delta_1 \langle A_{13} \rangle - iFg_1(x)\langle a \rangle - \Gamma_1 \langle A_{13} \rangle, \\ \langle \dot{A}_{23} \rangle &= i\Delta_2 \langle A_{23} \rangle - i(1 - F)g_2(x)\langle a \rangle - \Gamma_2 \langle A_{23} \rangle. \end{aligned} \quad (7)$$

With the following definitions

$$\mathbf{I}_\eta = \begin{pmatrix} \eta \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{Y} = \begin{pmatrix} a \\ A_{13} \\ A_{23} \end{pmatrix}$$

$$\mathbf{Z} = \begin{pmatrix} i\Delta_c - \kappa & -ig_1(x) & -ig_2(x) \\ -iFg_1(x) & i\Delta_1 - \Gamma_1 & 0 \\ -i(1 - F)g_2(x) & 0 & i\Delta_2 - \Gamma_2 \end{pmatrix} \quad (8)$$

the system of linear differential equations (7) can be written in a compact matrix notation

$$\langle \dot{Y} \rangle = \mathbf{Z} \langle Y \rangle + I_\eta. \quad (9)$$

The steady state solution of this system of linear differential equations is given by

$$\langle A_{13} \rangle_0 = \frac{ig_1(x)F\langle a \rangle}{(i\Delta_1 - \Gamma_1)}, \quad (10)$$

$$\langle A_{23} \rangle_0 = \frac{ig_2(x)(1-F)\langle a \rangle}{(i\Delta_2 - \Gamma_2)}. \quad (11)$$

Inserting Eqs. 11(a) and 11(b) into the Eq. 7 for the amplitude of the field we find

$$\begin{aligned} \langle \dot{a} \rangle = & (i\Delta_c - \kappa - F\gamma_1(x) - iFU_1(x) - (1-F)\gamma_2(x) \\ & - i(1-F)U_2(x))\langle a \rangle + \eta, \end{aligned} \quad (12)$$

where

$$\begin{aligned} U_1(x) = & \Delta_1 \frac{g_1^2(x)}{\Delta_1^2 + \Gamma_1^2}, U_2(x) = \Delta_2 \frac{g_2^2(x)}{\Delta_2^2 + \Gamma_2^2}, \\ \gamma_1(x) = & \Gamma_1 \frac{g_1^2(x)}{\Delta_1^2 + \Gamma_1^2}, \gamma_2(x) = \Gamma_2 \frac{g_2^2(x)}{\Delta_2^2 + \Gamma_2^2}, \end{aligned} \quad (13)$$

The dipole force acting on the rest Λ -type three-level atom is obtained,

$$\begin{aligned} \mathbf{F}(x) = & \dot{\mathbf{P}} = \frac{i}{\hbar} [H, \mathbf{P}] = -\hbar \nabla g_1(x) (a^\dagger A_{13} + A_{31} a) \\ & - \hbar \nabla g_2(x) (a^\dagger A_{23} + A_{32} a) \end{aligned} \quad (14)$$

and inserting Eqs. (11, 12) yields

$$\begin{aligned} \langle \mathbf{F}(x) \rangle_0 = & -\hbar \eta^2 \left(\frac{F(2+F)(\Delta_2^2 + \Gamma_2^2)\Delta_1(\nabla g_1(x))^2}{|det(\mathbf{Z})|^2} \right. \\ & \left. + \frac{(1-F)(3-F)(\Delta_1^2 + \Gamma_1^2)\Delta_2(\nabla g_2(x))^2}{|det(\mathbf{Z})|^2} \right), \end{aligned} \quad (15)$$

$det(\mathbf{Z})$ is the determinant of $\mathbf{Z}$, which is given by

$$\begin{aligned} det(\mathbf{Z}) = & (i\Delta_c - \kappa)(i\Delta_1 - \Gamma_1)(i\Delta_2 - \Gamma_2) \\ & + Fg_1^2(x)(i\Delta_2 - \Gamma_2) + (1-F)g_2^2(x)(i\Delta_1 - \Gamma_1). \end{aligned} \quad (16)$$

The expression is the dipole force. It is straightforward to find the expression for the friction force, the linear velocity dependence of the force for small velocities ($kv \ll \kappa$) due to the cavity dynamics. Use Eq. (12) in power of v to obtain the operator expectation values to first order [20],

$$\langle a \rangle_1 = \frac{v \cdot \nabla \langle a \rangle_0}{\Theta}, \quad (17)$$

where $\Theta = (i\Delta_c - \kappa - F\gamma_1(x) - iFU_1(x) - (1-F)\gamma_2(x) - i(1-F)U_2(x))\langle a \rangle$

Thus the expectation value of the force operator in first order of the velocity v ,

$$\begin{aligned} \langle \mathbf{F}(x) \rangle_1 = & -\hbar (\langle a^\dagger \rangle_0 \langle a \rangle_1 + \langle a^\dagger \rangle_1 \langle a \rangle_0 + \langle a \rangle_1 \langle a^\dagger \rangle_0 \\ & + \langle a \rangle_0 \langle a^\dagger \rangle_1) \nabla (F(1-F)U_1(x) \\ & + (1-F)(3-F)U_2(x)) \\ \equiv & -\beta v, \end{aligned} \quad (18)$$

β is called friction coefficient. Inserting Equation (17) into an expansion of the force Equation (18). Finally one obtains the friction force in the high-finesse optical cavity,

$$\begin{aligned} \langle F \rangle_1 = & -\frac{8\hbar v \eta^2 (\Delta_1^2 + \Gamma_1^2)(\Delta_2^2 + \Gamma_2^2)}{|det(\mathbf{Z})|^6} \cdot \\ & + (g_1(x)(\nabla g_1(x))(F(1-F)^2 A_{11} \\ & + 2F(1-F)B_{11} + FC_{11}) \\ & + g_2(x)(\nabla g_2(x))((1-F)F^2 A_{22} \\ & + 2(1-F)FB_{22} + (1-F)C_{22})) \cdot \\ & + (F(1-F)(\Delta_2^2 + \Gamma_2^2)\Delta_1 g_1(x)(\nabla g_1(x)) \\ & + (1-F)(3-F)(\Delta_1^2 + \Gamma_1^2)\Delta_2 g_2(x)(\nabla g_2(x))) \end{aligned} \quad (19)$$

where

$$\begin{aligned} A_{11} = & g_2^4(x)(\Delta_1^2 + \Gamma_1^2)(2\Gamma_2\Delta_1\Delta_2 + \Gamma_1(\Gamma_2^2 - \Delta_2^2)), \\ A_{22} = & g_1^4(x)(\Delta_2^2 + \Gamma_2^2)(2\Gamma_1\Delta_1\Delta_2 + \Gamma_2(\Gamma_1^2 - \Delta_1^2)), \\ B_{11} = & g_2^2(x)(\Delta_1^2 + \Gamma_1^2)(\Delta_2^2 + \Gamma_2^2)(Fg_1^2(x)\Gamma_2 \\ & + \kappa(\Delta_1\Delta_2 + \Gamma_1\Gamma_2) - \Delta_c(\Gamma_2\Delta_1 - \Gamma_1\Delta_2)), \\ B_{22} = & g_1^2(x)(\Delta_1^2 + \Gamma_1^2)(\Delta_2^2 + \Gamma_2^2)((1-F)g_2^2(x)\Gamma_1 \\ & + \kappa(\Delta_1\Delta_2 + \Gamma_1\Gamma_2) + \Delta_c(\Gamma_2\Delta_1 - \Gamma_1\Delta_2)), \\ C_{11} = & (\Delta_2^2 + \Gamma_2^2)^2(F^2g_1^4(x)\Gamma_1 + (2F\kappa g_1^2(x) \\ & - 2\kappa\Delta_1\Delta_c + \Gamma_1(\kappa^2 - \Delta_c^2))(\Delta_1^2 + \Gamma_1^2)), \\ C_{22} = & (\Delta_1^2 + \Gamma_1^2)^2((1-F)^2g_2^4(x)\Gamma_2 + (2(1-F)\kappa g_2^2(x) \\ & - 2\kappa\Delta_2\Delta_c + \Gamma_2(\kappa^2 - \Delta_c^2))(\Delta_2^2 + \Gamma_2^2)). \end{aligned} \quad (20)$$

The total force can be written as

$$\langle \mathbf{F}(x) \rangle \langle \mathbf{F}(x) \rangle_0 + \langle \mathbf{F}(x) \rangle_1 = \langle \mathbf{F}(x) \rangle_0 - \beta v \quad (21)$$

The equation (37) describes the total force of the Λ -type three-level atom within the high-finesse cavity, the first term is dipole force, which can be found from the steady state quantum average of the atom-cavity field interaction. The second term is the friction force. When the friction coefficient is positive, corresponding to cooling, while the friction coefficient is negative, leading to heating. The value of the friction coefficient can significantly modify the cooling process, so it is necessary to investigate the friction coefficient.

3 THE DIFFUSION COEFFICIENT AND THE EQUILIBRIUM TEMPERATURE

In the previous section we have obtained the friction coefficient, when $\beta > 0$, the atom can be cooled. However, momentum diffusion counteracts this cooling and prohibits that the atom completely stops at rest. In our paper, there are two major contributions to the momentum diffusion. The first referred to the momentum diffusion is due to the fluctuations of the dipole force. The other one is the random momentum transfer of absorbed and emitted photons.

We first calculate the momentum diffusion coefficient due to the fluctuations of the dipole force. The dipole force operator reads

$$F(x) = -\hbar \nabla g_1(x)(a^\dagger A_{13} + A_{31}a) - \hbar \nabla g_2(x)(a^\dagger A_{23} + A_{32}a). \quad (22)$$

For simplified calculation, we can derive equations for the expectation values of the normally ordered operator products, which yields

$$\langle \dot{\mathbf{X}} \rangle = (\Pi)_{9 \times 9} \langle \mathbf{X} \rangle + \eta \langle \Sigma \rangle, \quad (23)$$

with η denotes the pumping strengths of the external driving fields and $(\Pi)_{9 \times 9}$ is the coefficient matrix, follows

$$\Pi = \begin{bmatrix} (M_{11})_{3 \times 3} & (M_{12})_{3 \times 3} & (M_{13})_{3 \times 3} \\ (M_{21})_{3 \times 3} & (M_{22})_{3 \times 3} & (M_{23})_{3 \times 3} \\ (M_{31})_{3 \times 3} & (M_{32})_{3 \times 3} & (M_{33})_{3 \times 3} \end{bmatrix}, \quad (24)$$

where the several matrices $(M_{ij})_{3 \times 3}$ ($i, j = 1, 2, 3$) are given by

$$(M_{11})_{3 \times 3} = \begin{bmatrix} -(\kappa + \Gamma_1) & 0 & -(\Delta_1 - \Delta_c) \\ 0 & -(\kappa + \Gamma_2) & 0 \\ \Delta_1 - \Delta_c & 0 & -(\kappa + \Gamma_1) \end{bmatrix},$$

$$(M_{12})_{3 \times 3} = \begin{bmatrix} 0 & 0 & 0 \\ -(\Delta_2 - \Delta_c) & 0 & 0 \\ 0 & -2g_1(x)F & 2g_1(x) \end{bmatrix},$$

$$(M_{13})_{3 \times 3} = \begin{bmatrix} 0 & 0 & -ig_2(x) \\ 0 & 0 & ig_1(x) \\ 0 & g_2(x) & 0 \end{bmatrix},$$

$$(M_{21})_{3 \times 3} = \begin{bmatrix} 0 & \Delta_2 - \Delta_c & 0 \\ 0 & 0 & g_1(x) \\ 0 & 0 & -g_1(x)F \end{bmatrix},$$

$$(M_{22})_{3 \times 3} = \begin{bmatrix} -(\kappa + \Gamma_2) - 2g_2(x)(1 - F) & 0 & 0 \\ g_2(x) & -2\kappa & 0 \\ 0 & 0 & -2\Gamma_1 \end{bmatrix},$$

$$(M_{23})_{3 \times 3} = \begin{bmatrix} 2g_2(x) & g_1(x) & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

$$(M_{31})_{3 \times 3} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -g_2(x)(1 - F) \\ -ig_2(x)(1 - F) & ig_1(x)F & 0 \end{bmatrix},$$

$$(M_{32})_{3 \times 3} = \begin{bmatrix} -g_2(x)(1 - F) & 0 & 0 \\ -g_1(x)F & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

$$(M_{33})_{3 \times 3} = \begin{bmatrix} -2\Gamma_2 & 0 & 0 \\ 0 & -(\Gamma_1 + \Gamma_2) & -i(\Delta_1 - \Delta_2) \\ 0 & -i(\Delta_1 - \Delta_2) & -(\Gamma_1 + \Gamma_2) \end{bmatrix}. \quad (25)$$

and the variable X is defined as

$$\mathbf{X} \equiv \begin{pmatrix} X_1 \\ X_2 \\ X_3 \\ X_4 \\ X_5 \\ X_6 \\ X_7 \\ X_8 \\ X_9 \end{pmatrix} \equiv \begin{pmatrix} a^\dagger A_{13} + A_{31}a \\ a^\dagger A_{23} + A_{32}a \\ -i(a^\dagger A_{13} - A_{31}a) \\ -i(a^\dagger A_{23} - A_{32}a) \\ a^\dagger a \\ A_{31}A_{13} \\ A_{32}A_{23} \\ A_{31}A_{23} + A_{32}A_{13} \\ A_{31}A_{23} - A_{32}A_{13} \end{pmatrix} \quad (26)$$

Σ can be found as

$$\Sigma = \begin{pmatrix} A_{13} + A_{31} \\ A_{23} + A_{32} \\ -i(A_{13} - A_{31}) \\ -i(A_{23} - A_{32}) \\ a + a^\dagger \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (27)$$

According to the normally ordered operators, the dipole force can be expressed as

$$F(x) = -\hbar \nabla g_1(x)X_1 - \hbar \nabla g_2(x)X_2. \quad (28)$$

The part of the diffusion due to the dipole fluctuation can be written as

$$D_{dp} = \hbar^2 (\nabla \nabla g_1(x))^2 \text{Re} \int_0^\infty dt \langle \delta X_1(0) \delta X_1(t) \rangle + \hbar^2 (\nabla g_2(x))^2 \text{Re} \int_0^\infty dt \langle \delta X_2(0) \delta X_2(t) \rangle. \quad (29)$$

The Eq. (23) can be written as a homogeneous one for $\langle \tilde{X} \rangle = \langle X \rangle - X_0$, where X_0 is the steady-state value of $\langle X \rangle$. So we get

$$\frac{\partial \langle \tilde{X} \rangle}{\partial \tau} = (\Pi)_{9 \times 9} \langle \tilde{X} \rangle, \quad (30)$$

Due to the relation,

$$\langle \delta a_\nu \delta a_\mu \rangle = \langle a_\nu a_\mu \rangle - \langle a_\nu \rangle \langle a_\mu \rangle, \quad (31)$$

We can determine the correlation the correlation function $\langle \delta X_1(0) \delta X_1(t) \rangle$ and $\langle \delta X_2(0) \delta X_2(t) \rangle$ from the quantum regression theorem,

$$\frac{\partial \langle \tilde{X}(0) \tilde{X}(\tau) \rangle}{\partial \tau} = (\Pi)_{9 \times 9} \langle \tilde{X}(0) \tilde{X}(\tau) \rangle, \quad (32)$$

we obtain

$$\int_0^\infty \langle \delta X(0) \delta X(t) \rangle d\tau = -(\Pi)_{9 \times 9}^{-1} \langle \tilde{X}(0) \tilde{X}(0) \rangle \equiv \Xi, \quad (33)$$

The diffusion coefficient can be reduced to

$$D_{dp} = \hbar^2 (\nabla g_1(x))^2 \Xi_{11} + \hbar^2 (\nabla g_2(x))^2 \Xi_{22}. \quad (34)$$

The second contribution to the diffusion is the random momentum transfer of absorbed and emitted photons, the recoil contributes to the total diffusion by

$$\begin{aligned} D_s &= \hbar^2 k^2 \Gamma_1 \langle A_{31} A_{13} \rangle + \hbar^2 k^2 \Gamma_2 \langle A_{32} A_{23} \rangle \\ &= \hbar^2 k^2 \eta^2 \left(\frac{g_1^2(x) F^2 \Gamma_1 (\Delta_2^2 + \Gamma_2^2)}{|\det(\mathbf{Z})|^2} \right. \\ &\quad \left. + \frac{g_2^2(x) (1-F)^2 \Gamma_2 (\Delta_1^2 + \Gamma_1^2)}{|\det(\mathbf{Z})|^2} \right). \end{aligned} \quad (35)$$

The total diffusion can be obtained

$$D = D_{dp} + D_s. \quad (36)$$

Hence, we can obtain the equilibrium temperature,

$$k_B T = \frac{D}{\beta}. \quad (37)$$

Using the Heisenberg equations and the quantum regression theorem we obtain the full analytical expressions for the friction, diffusion coefficients and the equilibrium temperature. Because we have many more parameters available to control the cooling process in the Λ -type three-level atomic system, it is necessary to study the variation of the friction, diffusion coefficients and the equilibrium temperature with these parameters.

4 Numerical calculation

Using the theory described in the previous paragraphs, we have given a detailed analysis of the cooling process of the Λ -type three-level atom in the high-finesse optical cavity. For simplicity, we will only consider one-dimensional situation, and assume $K_1 = K_2 = k$, $\Gamma_1 = \Gamma_2 = \Gamma$ in the following. Our results are represented in scaled quantities: the parameters Δ_c , Δ_1 , Δ_2 , g_0 , Γ and η are divided by κ . The friction, diffusion coefficients and the equilibrium temperature are calculated numerically and shown graphically. For a suitable choice of the controllable parameters, the equilibrium temperature can be cool down below the Doppler limit. Fig. 2 shows that the average of friction, diffusion coefficients and the equilibrium temperature along the cavity axis over a length of one wavelength are plotted as functions of the cavity detuning Δ_c . The parameters are set to $g_0 = 8\kappa$, $\Delta_1 = 8\kappa$, $\Delta_2 = 7\kappa$, $\Gamma = 1.4\kappa$ and $\eta = 1.5\kappa$. As is well known, for cooling a single atom in a free standing wave, the attainable temperature is given by $T_D = \hbar\gamma/k_B$, which is the so-called Doppler limit. Here γ represents the nature linewidth. In our case of cavity




Fig. 2. The average of the friction, diffusion coefficients and the equilibrium temperature along the cavity axis over a length of one wavelength are plotted as functions of the cavity detuning Δ_c . (a) the average of the friction coefficient, (b) the average of the diffusion coefficient, (c) the equilibrium temperature. The parameters are set to $g_0 = 8\kappa$, $\Delta_1 = 8\kappa$, $\Delta_2 = 7\kappa$, $\Gamma = 1.4\kappa$ and $\eta = 1.5\kappa$.

cooling of a Λ -type three-level atom, the equilibrium temperature is proportion to $\hbar\kappa/k_B$, κ expresses the cavity loss rate. If κ is smaller than the γ , we can obtain the final temperatures lower than the Doppler limit, which can be seen from the Fig. 2(c).

For a Λ -type three-level atom in a high-finesse cavity, there are more parameters available to control the friction, diffusion coefficients and the equilibrium temperature, it is necessary to investigate the variation of the cooling process with these parameters. To select detunings which are experimentally advantageous for cavity cooling and trapping of the Λ -type three-level atom, it is necessary to investigate the average of the friction and diffusion coefficients that the atom will experience in the high-finesse optical cavity. Figure 3 shows that the average of the friction and diffusion coefficients along the cavity axis over the length of λ are plotted as functions of atomic detunings Δ_1 and Δ_2 for $\Delta_c = 0$. The other parameters are set to $g_0 = 8\kappa$, $\Gamma = 1.4\kappa$ and $\eta = 1.5\kappa$. It can be found that the atomic detunings affect the average of the friction and




Fig. 3. (a) the average of the friction coefficient, (b) the average of the diffusion coefficient (c) the equilibrium temperature along the cavity axis over a length of λ are plotted as functions of atomic detunings Δ_1 and Δ_2 for $\Delta_c = 0$. The other parameters are set to $g_0 = 8\kappa$, $\Gamma = 1.4\kappa$ and $\eta = 1.5\kappa$.




Fig. 4. The dependence of (a) the average of friction, (b) the average of diffusion coefficients and (c) the equilibrium temperature on the coupling strengths $g_0 = 4\kappa$, $g_0 = 8\kappa$, $g_0 = 12\kappa$. The other parameters are set to $\Delta_1 = 8\kappa$, $\Delta_2 = 7\kappa$, $\eta = 1.5\kappa$, $\Gamma = 1.4\kappa$.

diffusion coefficients significantly. For a suitable choice of the parameters, we can obtain the low equilibrium temperature, which can be seen from the Fig. 3(c). According to Fig. 3, we give a especial case for a fixed values of Δ_1 and Δ_2 . On one hand, we consider the dependence of the average of friction, diffusion coefficients and equilibrium temperature on the coupling strengths, on the other hand, we investigate the average of friction, diffusion coefficients and equilibrium temperature with the pumping strengths.

Firstly, we demonstrate the dependence of the average of friction, diffusion coefficients and equilibrium temperature on the coupling strengths $g_0 = 4\kappa$, $g_0 = 8\kappa$, $g_0 = 12\kappa$ in Fig. 4. The other parameters are set to $\Delta_1 = 8\kappa$, $\Delta_2 = 7\kappa$, $\eta = 1.5\kappa$, $\Gamma = 1.4\kappa$. It can found that the friction and diffusion coefficients are enhanced, while the equilibrium temperature is reduced with increase of the coupling strengths. As is well known, in the strongly coupled regime, a strong friction force exists, there is also a strong heating rate. For a batter understanding of the strong friction force, the interpretation can use the dressed states and sisyphus cooling mechanism. For the Λ -type three-level atom coupled with the cavity, the eigenvalues corre-




Fig. 5. Energies of the Λ -type three-level atom within cavity system (dressed states) as a function of the axial position for (a) $g_0 = 12\kappa$, (b) $g_0 = 8\kappa$.

spond to the energy are E_1 , E_2 , and E_3 respectively ($\hbar \equiv 1$).

$$\begin{aligned}
 E_1 &= \omega_p - \frac{(2\Delta_2 - \Delta_1)}{3} \\
 &\quad + \sqrt[3]{q - \sqrt{q^2 + p^3}} + \sqrt[3]{q + \sqrt{q^2 + p^3}}, \\
 E_2 &= \omega_p - \frac{(2\Delta_2 - \Delta_1)}{3} \\
 &\quad - \frac{1}{2}(\sqrt[3]{q - \sqrt{q^2 + p^3}} + \sqrt[3]{q + \sqrt{q^2 + p^3}}) \\
 &\quad - \frac{\sqrt{-3}}{2}(\sqrt[3]{q - \sqrt{q^2 + p^3}} - \sqrt[3]{q + \sqrt{q^2 + p^3}}), \\
 E_3 &= \omega_p - \frac{(2\Delta_2 - \Delta_1)}{3} \\
 &\quad - \frac{1}{2}(\sqrt[3]{q - \sqrt{q^2 + p^3}} + \sqrt[3]{q + \sqrt{q^2 + p^3}}) \\
 &\quad + \frac{\sqrt{-3}}{2}(\sqrt[3]{q - \sqrt{q^2 + p^3}} - \sqrt[3]{q + \sqrt{q^2 + p^3}}), \quad (38)
 \end{aligned}$$

Where

$$\begin{aligned}
 p &= -\left(\frac{2\Delta_2 - \Delta_1}{3}\right)^2 - \frac{g_1^2(x) + g_2^2(x) + 4\Delta_2(\Delta_2 - \Delta_1)}{12}, \\
 q &= \left(\frac{2\Delta_2 - \Delta_1}{3}\right)^3
 \end{aligned}$$




Fig. 6. The dependence of (a) the average of friction, (b) the average of diffusion coefficients and (c) the equilibrium temperature on the pumping strengths $\eta = 0.5\kappa$, $\eta = 1.0\kappa$, $\eta = 1.5\kappa$. The other parameters are set to $\Delta_1 = 8\kappa$, $\Delta_2 = 7\kappa$, $g_0 = 8\kappa$, $\Gamma = 1.4\kappa$.

$$\begin{aligned}
 &+ \frac{(2\Delta_2 - \Delta_1)(g_1^2(x) + g_2^2(x) + 4\Delta_2(\Delta_2 - \Delta_1))}{24} \\
 &+ \frac{g_1^2(x)(\Delta_2 - \Delta_1)}{8}, \quad (39)
 \end{aligned}$$

here $g_1(x)$ and $g_2(x)$ are the coupling strength, $\Delta_1 = \omega_P - \omega_{31}$ and $\Delta_2 = \omega_P - \omega_{32}$ are the atomic detunings. The ground state energy is E_0 . Eigenvalues E_1 , E_2 , and E_3 as a function of axial position are shown in Fig. 5. In this atom-cavity system, cavity cooling arise when the system is excited near a minimum of one of the dressed states, the system will remain in this state while the atom moves further and thus the system energy will vary according to corresponding eigenvalues. For the Λ -type three-level atom system, after being excited in the potential valley of the dressed state, the stronger of the coupling strength, the steeper hills for the Λ -type three-level atom has to climb, which can be seen in Fig.5(a) and in Fig.5(b). Hence, friction coefficient become stronger with the bigger coupling strength. On the contrary, the momentum diffusion can be affected by the fluctuations of atomic dipole coupled to the cavity field, because the Λ -type three-level atom ex-

perience a stronger force with the increase of the coupling strength, momentum diffusion coefficient will be enhanced by increase of the coupling strength. However, for a suitable choice of parameters, the atomic cooling temperature can be down to lower than the Doppler temperature limit T_D , which can be seen from the Fig. 4(c).

Moreover, we illustrate the dependence of the average of friction, diffusion coefficients and equilibrium temperature on the pumping field strengths $\eta = 0.5\kappa, \eta = 1.0\kappa, \eta = 1.5\kappa$ in Fig. 6. The other parameters are set to $\Delta_1 = 8\kappa, \Delta_2 = 7\kappa, g_0 = 8\kappa, \Gamma = 1.4\kappa$. It can be found that with the increase of the pumping strengths, the friction and diffusion coefficients are enhanced, while the equilibrium temperature is reduced. This is due to the friction and diffusion coefficients are proportion to quadratic of the pumping field, the average of friction and diffusion coefficients can be affected remarkably by the increase of the pumping field strength. While the equilibrium temperature is proportion to $\hbar\kappa/k_B$, we can obtain the final temperature lower than the Doppler temperature limit.

5 CONCLUSION

In this paper, using the master equation and the Heisenberg equations, we have derived the analytical solution of the dipole force and friction force for the Λ -type three-level atom in the high-finesse cavity. The friction coefficient can significantly affect the cooling process, however, momentum diffusion counteracts this cooling and prohibit that the atom completely stops at rest. In our paper, there are two major contributions to the momentum diffusion. One is the random momentum transfer of absorbed and emitted photons, the second referred to the momentum diffusion is due to the fluctuations of the dipole force. According to quantum regression theorem, we obtain the diffusion coefficient and the equilibrium temperature. The friction, diffusion coefficients and the equilibrium temperature are calculated numerically and shown graphically as a function of controlling parameters. For a suitable choice of parameters, the atomic cooling temperature can be down to lower than the Doppler temperature limit.

This work was partly supported by the National Natural Science Foundation of China under Grant No. 10704031, the National Natural Science Foundation of China for Fostering Talents in Basic Research under Grant No. J0630314, the Fundamental Research Fund for Physics and Mathematics of Lanzhou University under Grant No. Lzu05001, and the Nature Science Foundation of Gansu under Grant No. 3ZS061 – A25 – 035.

References

1. P. Horak, G. Hechenblaikner, K. M. Gheri, H. Stecher and H. Ritsch. Phys. Rev. Lett. **79** 4974 (1997).
2. P. Maunz, T. Puppe, I. Schuster, N. Syassen, P.W. H. Pleskes and G. Rempe. Nature (London). **428** 50 (2004).
3. S. Chu. Rev. Mod. Phys. **70** 685 (1998).

4. C. N. Cohen-Tannoudji. Rev. Mod. Phys. **70** 707 (1998).
5. W. D. Phillips. Rev. Mod. Phys. **70** 721 (1998).
6. M. Viteau, A. Chotia, M. Allegrini, N. Bouloufa, O. Dulieu, D. Comparat and P. Pillet. Science **321** 232 (2008).
7. J. G. Danzl, E. Haller, M. Gustavsson, M. j. Mark, R. Hart, N. Bouloufs, O. Dulieu, H. Ritsch and H. C. Nägerl. Science **321** 1062 (2008).
8. M. Y. Vilensky, Y. Prior, and I. S. Averbukh, Phys. Rev. Lett. **99** 103002 (2007).
9. M. Hijkema and B. Weber. Nature Phys. **3** 253 (2007).
10. T. Wilk and S. C. Webster. Science. **317** 488 (2007).
11. F. Brennecke, T. Donner, S. Ritter, T. Bourdel, M. Kihl and T. Esslinger. Nature. **450** 268 (2007).
12. J. M. Zhang, F. C. Cui, D. L. Zhou and W. M. Liu. Phys. Rev. A **79** 033401 (2009).
13. G. Szirmai, D. Nagy and P. Domokos. Phys. Rev. Lett. **102** 080401 (2009).
14. J. H. Wu, M. Artoni and G. C. La Rocca. Phys. Rev. Lett. **103** 133601 (2009).
15. H. J. Metcalf, P. V. Straten 1999 Laser cooling and trapping (Springer-Verige New York) P57.
16. S. Nußmann, K. Murr, M. Hijkema, B. Weber, A. Kuhn, G. Rempe. Nature phys **1** 122 (2005).
17. S. Zippilli, G. Morigi. Phys. Rev. Lett. **72** 143001 (2005).
18. S. Zippilli, G. Morigi. Phys. Rev. A. **72** 053408 (2005).
19. T. Salzbergur, P. Domokos, and H. Ritsch. Phys. Rev. A. **72** 033805 (2005).
20. G. Hechenblaikner, M. Gangl, P. Horak, and H. Ritsch. Phys. Rev. A. **58** 3030 (1998).
21. P. R. Hemmer and M. G. Prentss. J. Opt. Soc. Am. B. **5** 1613 (1998).
22. L. W. Liu, L. Tan, and Y. Xu. J. Mod. Opt **56** 968(2008)
23. A. E. Siegman. Laser; University Sciene Books: Mill valley (1986) P502.
24. I. Protsenko, P. Domokos, V. Lefevre, J. Hare, J. M. Rainond, and L. Davidovich. Phys. Rev. A **59** 1667 (1999)
25. H. M. Wiseman, and G. J. Milburn. Phys. Rev. A. **47** 642(1993).

